Spherical Solutions in Einstein-Aether Theory: Static Aether and Stars

Christopher Eling* and Ted Jacobson†

Department of Physics, University of Maryland

College Park, MD 20742-4111 USA

arXiv:gr-qc/0603058v4 6 Dec 2009

Abstract

The time independent spherically symmetric solutions of General Relativity (GR) coupled to a dynamical unit timelike vector are studied. We find there is a three-parameter family of solutions with this symmetry. Imposing asymptotic flatness restricts to two parameters, and requiring that the aether be aligned with the timelike Killing field further restricts to one parameter, the total mass. These "static aether" solutions are given analytically up to solution of a transcendental equation. The positive mass solutions have spatial geometry with a minimal area 2-sphere, inside of which the area diverges at a curvature singularity occurring at an extremal Killing horizon that lies at a finite affine parameter along a radial null geodesic. Regular perfect fluid star solutions are shown to exist with static aether exteriors, and the range of stability for constant density stars is identified.

*Electronic address: cteling@physics.umd.edu

†Electronic address: jacobson@umd.edu

I. INTRODUCTION

For nearly a century Lorentz invariance has been a basic assumption in theoretical physics. However, various approaches to the problem of quantum gravity question whether Lorentz symmetry is truly fundamental. In this context, it is useful to consider low energy Lorentz violating (LV) models from the point of view of effective field theory. It makes sense to distinguish LV effects in the matter sector from LV effects in the much more weakly coupled (at low energies) gravitational sector. LV effects in the matter sector are tightly constrained by high precision tests of Lorentz invariance [1]. Lorentz violation in the gravitational sector was studied by Will and Nordvedt [2] in the early 1970's, and it was pursued in the 1980's by Gasperini [3] motivated by singularity prevention, and by Kostelecky and Samuel [4] motivated by the possibility of spontaneous LV in string theory. More recently a number of approaches to incorporating LV into gravity have been explored; see for example Refs. [5, 6, 7, 8, 9, 10, 11] and references therein.

Here we study the theory of a unit timelike vector field u^a coupled only to gravity, as an LV modification of General Relativity (GR). To preserve general covariance this field must be dynamical. It can be thought of as the 4-velocity of a preferred frame, which breaks boost symmetry at every point of spacetime. This property it has in common with the 19th century concept of the aether, and like that aether it is also a dynamical system. Hence a suitable name for it is the "aether", although it has nothing to do with electromagnetism. Since the aether is coupled to Einstein GR we refer to the theory as "Einstein-Aether" theory; we sometimes use the nickname "ae-theory" for short. For a review of the motivation, history and recent status of the Einstein-Aether theory, see [12] and the references therein.

Observational constraints on the Einstein-Aether theory have been determined from PPN analysis [13, 14, 15], stability and energy positivity [16, 17, 18, 19], primordial nucleosynthesis [20], and Cerenkov radiation [18]. The combined constraints from all of these are reviewed in Ref. [15], and constraints from radiation damping are discussed in [21]. Also important for determining theoretical viability and observational constraints are the properties the spherically symmetric static solutions. These solutions were previously obtained in the the special case where the aether dynamics is Maxwell-like [4, 6]. It was shown in [6] that the Reissner-Nordstrom metric in a spherically symmetric static gauge with fixed norm is a solution, and it was claimed (incorrectly, as shown here) that this is the only solution

in that special case. The asymptotic weak field limit of the general case was studied in Ref. [13], where it was found that there is a two-parameter family of asymptotically flat spherical, static solutions. A thorough examination of the fully nonlinear solutions has not been carried out before.

In this paper and its companion (devoted to black holes) [22] we complete a general survey of the time-independent spherically symmetric solutions. In section II of this paper the action and field equations for the Einstein-Aether theory are reviewed. Specializing to stationary, spherical symmetry, section III studies the structure of the field equations as ordinary differential equations and shows that locally there is a three parameter family of vacuum solutions. Imposing asymptotic flatness reduces the number of parameters to two.

In section IV the general solution in which the aether vector is aligned with the timelike Killing field is found. Being a unit vector, the aether is completely determined by the metric in this case. This solution is asymptotically flat and described by one free total mass parameter. For negative total mass there is a naked singularity at the origin. The positive mass solution has a wormhole-like spatial geometry, reaching a minimum area 2-sphere at some radius like the Schwarzschild solution. Unlike in the Schwarzschild solution this throat is not on a horizon. Inside the throat the spheres re-expand to infinitely large size in finite affine parameter along a radial null geodesic and finite or infinite proper distance depending on the coupling parameters in the Lagrangian. When the distance is finite the internal infinity is singular, and it occurs at a would-be extremal Killing horizon. When the distance is infinite the metric is asymptotically singular.

In section V it is first shown that pure aether stars do not exist, i.e. there are no asymptotically flat self-gravitating aether solitons with a regular origin. (Given the results of the previous section, regular aether stars could only possibly have existed if the aether were not everywhere aligned with the Killing vector.) Next it is shown that in the presence of a perfect fluid, regular asymptotically flat star solutions exist and are parameterized (for a given equation of state) by the central pressure. For the case of constant density the star solutions are found by matching numerical integration for the interior to the vacuum solution found in section IV. As in GR, for a given density there is a maximum mass. Utilizing the critical behavior of the mass as a function of stellar radius R it is shown that if they are stable at small mass, these stars are unstable beyond the maximum mass. We conclude in section VI with a discussion of open questions raised by these results.

II. EINSTEIN-AETHER ACTION

The action for Einstein-Aether theory is the most general diffeomorphism invariant functional of the spacetime metric g_{ab} and aether field u^a involving no more than two derivatives,

$$S = \frac{1}{16\pi G}\int \sqrt{-g}\; L\; d^4x \tag{1}$$

where

$$L = -R - K^{ab}{}_{mn}\nabla_a u^m \nabla_b u^n - \lambda(g_{ab}u^a u^b - 1). \tag{2}$$

Here R is the Ricci scalar, $K^{ab}{}_{mn}$ is defined as

$$K^{ab}{}_{mn} = c_1 g^{ab}g_{mn} + c_2\delta^a_m\delta^b_n + c_3\delta^a_n\delta^b_m + c_4 u^a u^b g_{mn} \tag{3}$$

where the c_i are dimensionless constants, and λ is a Lagrange multiplier enforcing the unit timelike constraint. This constraint restricts variations of the aether to be spacelike, hence ghosts need not arise. A term of the form $R_{ab}u^a u^b$ is not explicitly included as it is proportional to the difference of the c_2 and c_3 terms in (1) via integration by parts. The metric signature is $(+---)$ and the units are chosen so that the speed of light defined by the metric g_{ab} is unity. In spherical symmetry the aether is hypersurface orthogonal, hence it has vanishing twist $\omega_a = \epsilon_{abcd}u^b\nabla^c u^d$. When u^a is a unit vector the square of the twist is a combination of the c_1, c_3, and c_4 terms in the action (1),

$$\omega_a\omega^a = -(\nabla_a u_b)(\nabla^a u^b) + (\nabla_a u_b)(\nabla^b u^a) + (u^b\nabla_b u_a)(u^c\nabla_c u^a). \tag{4}$$

The c_4 term can thus be absorbed by making the replacements $c_1 \to c_1 + c_4$ and $c_3 \to c_3 - c_4$.

Following the observational constraints we assume here when studying the fluid star solutions that the only significant coupling of u^a to matter is through a universal "matter metric" $g^{\rm matter}_{ab} = g_{ab} + \sigma u_a u_b$, where σ is a constant. Replacing g_{ab} by $g^{\rm matter}_{ab}$ as the independent metric field in the action returns an action with the same form as (1) but with new values of the constants $c_{1,2,3,4}$ that depend on σ [23]. Hereafter we assume that such a field redefinition has already been performed, so that g_{ab} is the metric to which matter couples universally. The absence of any other coupling of u^a to matter has no theoretical justification in this purely phenomenological approach, and may be regarded as unnatural. However our goal here is just to explore consequences of gravitational Lorentz violation in a phenomenologically viable setting. It remains an open question whether this can emerge as an approximation to a more fundamental underlying theory.

The field equations from varying (1) plus a matter action (coupled only to the metric) with respect to g^{ab}, u^a and λ are given by

$$G_{ab} = T^{(u)}_{ab} + 8\pi G T^{M}{}_{ab} \tag{5}$$

$$\nabla_a J^a{}_m - c_4 \dot{u}_a \nabla_m u^a = \lambda u_m, \tag{6}$$

$$g_{ab} u^a u^b = 1, \tag{7}$$

where

$$J^a{}_m = K^{ab}{}_{mn} \nabla_b u^n. \tag{8}$$

The aether stress tensor is given by [13]

$$\begin{aligned} T^{(u)}{}_{ab} = {} & \nabla_m (J_{(a}{}^m u_{b)} - J^m{}_{(a} u_{b)} - J_{(ab)} u^m) \\ & + c_1 \left[(\nabla_m u_a)(\nabla^m u_b) - (\nabla_a u_m)(\nabla_b u^m) \right] \\ & + c_4 \, \dot{u}_a \dot{u}_b \\ & + \left[u_n (\nabla_m J^{mn}) - c_4 \dot{u}^2 \right] u_a u_b \\ & - \frac{1}{2} L_u g_{ab}, \end{aligned} \tag{9}$$

where $L_u = -K^{ab}{}_{mn} \nabla_a u^m \nabla_b u^n$. The Lagrange multiplier λ has been eliminated from (9) by solving for it via the contraction of the aether field equation (6) with u^a.

Some words about terminology are in order. Spacetimes admitting a timelike Killing vector field ξ^a are generally called *stationary*. In the special case where ξ^a is hypersurface orthogonal, and therefore invariant under a time reflection $t \to -t$, the spacetime is said to be *static*. A stationary aether field u^a on a stationary spacetime is one whose Lie derivative with respect to ξ^a vanishes. If the spacetime is static, one might be tempted to say the aether is "static", however this is not really appropriate since the aether itself breaks the Killing time reflection symmetry. The solutions studied in this paper involve a static metric coupled to a stationary aether. This general situation will be called here "stationary spherical symmetry". An important special case occurs when the aether is parallel to the Killing vector. We refer to this special case as "static aether". Such an aether changes sign under the Killing time reflection, however the action (1) is invariant under $u^a \to -u^a$ so the sign of u^a has no physical meaning. Note that regular black holes cannot have static aether fields since the Killing vector is null, not timelike on the horizon.

III. CLASSIFICATION OF STATIONARY SPHERICAL SOLUTIONS

Stationary spherically symmetric solutions describe, for example, a black hole or the exterior of a time-independent star. In spherical symmetry all stationary metrics are static [24]. The line element can be written in Schwarzschild type coordinates,

$$ds^2 = e^{A(r)}dt^2 - B(r)dr^2 - r^2 d\Omega^2, \tag{10}$$

and the aether field takes the form

$$u = a(r)\partial_t + b(r)\partial_r. \tag{11}$$

The unit constraint on u^a becomes

$$e^{A(r)}a(r)^2 - B(r)b(r)^2 = 1, \tag{12}$$

which can be used for example to eliminate $b(r)$. The t-component of the aether field equation (6) can be used to solve for λ in this case, and the remaining field equations reduce to five ODE's: the tt, rr, tr, and $\theta\theta$ components of the metric field equation and the r component of the aether field equation. These five equations involve the eight functions $\{A'', A', A, B', B, a'', a', a\}$, where prime denotes differentiation with respect to the argument r which is suppressed. The equations are too complicated to be worth writing down here, so we shall just describe their structure. Using the tt and $\theta\theta$ metric equations along with the r component of the aether equation, one can solve for A'', a'', and B' in terms of the remaining five functions $\{A', A, B, a', a\}$. It turns out that only one additional piece of information remains in the tr and rr equations, which can be used to solve (for example) for B in terms of $\{A', A, a', a\}$. Finally, $A(r_0)$ at any given value $r = r_0$ can be chosen at will by allowing for an appropriate scaling of the t coordinate. At a given r_0 value, the remaining three values $\{A'(r_0), a'(r_0), a(r_0)\}$ then determine a (local) solution by integration with respect to r. This shows that there is in general a three-parameter family of spherically symmetric stationary solutions.

To illustrate this reasoning in a more familiar setting, we can apply it to the field equations of pure GR in Schwarzschild coordinates,

$$G_{tt} \propto rB' - B + B^2 = 0 \tag{13}$$

$$G_{rr} \propto rA' - B + 1 = 0 \tag{14}$$

$$G_{\theta\theta} \propto 2rA''B + A'(2B - rB') + rA'^2B - 2B' = 0. \tag{15}$$

These can be used to solve for B', B, and A'' in terms of A' and A. Using the freedom to scale t the initial value $A(r_0)$ can be fixed at will, so we recover the well-known fact that static spherically symmetric solutions in GR are characterized by one free parameter, in this case the value of $A'(r_0)$. In the Einstein-Aether theory the aether vector and its derivative provide two additional degrees of freedom at each point.

Birkhoff's theorem in GR states that the only spherically symmetric solution is static and given (up to coordinate freedom) by the Schwarzschild metric. The radial tilt of the aether provides another local degree of freedom in ae-theory, so spherical solutions need not be time-independent. But, as we have seen, even restricting to stationary spherically symmetry ae-theory has more solutions. In this paper we will focus primarily on the static aether solutions, which form a one-parameter family. Black hole solutions, which comprise a different family, are studied in a companion paper.

Numerical integration of the ae-theory field equations as ODE's out from some arbitrary point r_0 with generic initial conditions yields singularities in $A(r)$, $B(r)$, and $a(r)$. However, there is a two-parameter family of asymptotically flat solutions. This was first found in Ref. [13] using a perturbative expansion about infinity. Asymptotic flatness was imposed there by assuming regular power series expansions about $x = 1/\rho = 0$, where ρ is the isotropic radial coordinate. Asymptotic flatness can also be imposed using the "shooting method". This is simple to implement here since it is only necessary to tune one of the three initial values $\{A'(r_0), a'(r_0), a(r_0)\}$ so that, for example, $A(r)$ approaches a constant value as $r \to \infty$. The field equations then automatically enforce the remaining asymptotic flatness conditions. In GR, by contrast, asymptotic flatness is a consequence of the vacuum field equations without any tuning of initial data, so the one-parameter family of local (Schwarzschild) solutions is automatically asymptotically flat.

IV. STATIC AETHER

In this section we obtain the static aether solution, where the aether vector u^a is proportional to the timelike Killing field ξ^a and therefore entirely determined by the metric.

A. Field equations with static aether

Using the Schwarzschild type coordinates in (10) and (11), the static aether has $b(r) = 0$ and $a(r) = \exp(-A(r)/2)$, i.e.

$$u = e^{-A/2}\partial_t. \tag{16}$$

In this case the c_2 and c_3 terms drop out of the field equations. To see why, note that (16) implies $\nabla_a u^a = 0$, so all variations of c_2 term in the action (1) vanish. In addition, the normalization $u_a u^a = 1$ implies $u_b \nabla_a u^b = 0$. These conditions together with spherical symmetry imply that the derivative of u^b has the form

$$\nabla_a u^b = u_a s^b, \tag{17}$$

where s^b is a radial vector orthogonal to u^a. (We note in passing that contraction of (17) with u^a reveals that s^b is the acceleration of the aether worldlines.) Therefore $(\nabla_a u^b)(\nabla_b u^a)$ vanishes, so the variation of the volume element in the c_3 term of the action (1) vanishes. The remaining variation of the c_3 term is proportional to $u_a s^b \delta(\nabla_b u^a) = s^b \delta(u_a \nabla_b u^a)$, which vanishes for all variations $(\delta g_{ab}, \delta u^a)$ preserving the normalization $g_{ab} u^a u^b = 1$. Moreover, as explained in Section II, in spherical symmetry the c_4 term can be absorbed into the c_1 and c_3 terms, hence the solutions with static aether are fully characterized by the case with only c_1 non-zero.

When only c_1 is nonzero the aether field equation (6) reduces to

$$c_1 \nabla^a \nabla_a u^b = \lambda u^b. \tag{18}$$

Using (17) this becomes

$$c_1 u^a \nabla_a s^b = \lambda u^b. \tag{19}$$

Contraction of the left hand side of (19) with s_b is proportional to $u^a \nabla_a s^2$, which vanishes since s^2 is a scalar that must be constant along the Killing direction parallel to u^a. Therefore both sides are parallel to u^b, so the aether equation only determines λ. Contracting both sides of (19) with u_b we find

$$\lambda = -c_1 s^2, \tag{20}$$

having made use of (17) and $u^b \nabla_a s_b = -s^b \nabla_a u_b$, which follows from $u^b s_b = 0$.

The metric field equation is $E_{ab} = G_{ab} - T^u_{ab} = 0$, and the tt, rr, and $\theta\theta$ components of E_{ab} are given by

$$E_{tt} = (e^A/r^2B)\Big[(-1 + B + rB'/B) - \nu(8rA' + r^2A'^2 - 2r^2A'B'/B + 4r^2A'')\Big] \quad (21)$$

$$E_{rr} = r^{-2}(1 - B + rA' + \nu r^2A'^2) \quad (22)$$

$$E_{\theta\theta} = B^{-1}\Big[(2rA' - 2rB'/B + r^2A'^2 - r^2A'B'/B + 2r^2A'')/4 - \nu r^2A'^2\Big], \quad (23)$$

where for notational convenience we have introduced the symbol

$$\nu = \frac{c_1}{8}. \quad (24)$$

Using the E_{rr} equation one can solve for B,

$$B = 1 + rA' + \nu\, r^2A'^2, \quad (25)$$

Substituting this solution for B into the E_{tt} and E_{rr} equations, we find that the equations are redundant and the system is described by the second order ODE

$$r^2A'' + 2rA' + r^2A'^2 + \nu\, r^3A'^3 = 0. \quad (26)$$

A constant shift of A can be absorbed by a scaling of the t coordinate, hence there is just a one parameter family of solutions. As in GR, the solutions in this family are all asymptotically flat.

B. Static aether solutions: general analysis

To solve (26) we define the function $Y(r)$ by

$$Y = rA', \quad (27)$$

in terms of which the solution for B becomes

$$B = 1 + Y + \nu Y^2 \quad (28)$$

and Eqn. (26) for A becomes

$$dY/dr = -(Y/r)(1 + Y + \nu Y^2) \quad (29)$$

The problem is thus reduced to quadratures: integration of this equation yields $Y(r)$, which also directly yields B via (28). To determine A we combine (27) and (29) to obtain

$$dA/dY = -1/(1 + Y + \nu Y^2), \tag{30}$$

which yields $A(Y)$ by integration.

The character of the solutions is evidently controlled by the roots of B,

$$Y_\pm = (-1 \pm \sqrt{1 - 4\nu})/(2\nu), \tag{31}$$

in terms of which we have

$$B = \nu(Y - Y_-)(Y - Y_+). \tag{32}$$

The nature of the roots depends on the value of ν. We consider here only positive ν, since that is required by positivity of the energy of linearized spin-0 waves [19], and we restrict to $\nu < 1/4$ since the Newton constant $G_{\rm N} = G/(1 - c_{14}/2)$ becomes negative beyond this limit. In the pure c_1 case also the stability [16] or positive energy [19] of linearized waves requires $c_1 < 1$, or $\nu < 1/8$. One can visualize the roots graphically: they are the intersections of the line $Y + 1$ with the inverted parabola $-\nu Y^2$. When $\nu = 1/4$ the parabola is tangent to the line, and the two roots coincide at $Y = -2$. The larger root approaches -1 as $\nu \to 0$, while the smaller root approaches $-\infty$, hence in the range $0 \le \nu < 1/4$ the roots fall within the ranges

$$-\infty \le Y_- < -2, \qquad -2 < Y_+ \le -1. \tag{33}$$

Note that $Y_- = 1/(\nu Y_+)$, and $\nu = -(1 + Y_+)/Y_+^2$.

We can integrate (30) and (29) to find both A and Y using the factorization (32) and partial fractions. The result is

$$N = e^A = \left(\frac{1 - Y/Y_-}{1 - Y/Y_+}\right)^{\frac{-Y_+}{2+Y_+}} \tag{34}$$

and

$$\frac{r_{\rm min}}{r} = \left(\frac{Y}{Y - Y_-}\right)\left(\frac{Y - Y_-}{Y - Y_+}\right)^{\frac{1}{2+Y_+}}, \tag{35}$$

where $r_{\rm min}$ is an integration constant. The graph of $r/r_{\rm min}$ vs. Y is plotted in Fig. 1, for the case $c_1 = 1$. The values of the sphere radius r and metric functions B and N at the special values of Y are given in Table I.

FIG. 1: Graph of $r/r_{\min}$ vs. Y for $c_1 = 1$. The curves approach 1 asymptotically on both sides. The range $(Y_+, 0)$ defines a negative mass solution with naked singularity at $Y = Y_+$ and asymptotically flat region at $Y \to 0$. The range $(0, \infty)$ defines a positive mass solution, with a minimal 2-sphere as $Y \to \infty$. The range $(-\infty, Y_-)$ continues that solution to the other side of the minimal sphere, with a singularity at a sphere of infinite radius at $Y = Y_-$. This sphere lies at finite radial distance if $c_1 < 3/2$. There is no solution with timelike aether in the range (Y_-, Y_+) since the radial coordinate is timelike there (see section IV B 4).

TABLE I: Sphere radius and metric functions at special Y values.

Y	Y_-	Y_+	0	$\pm\infty$
r	∞	0	∞	$r_{\min}$
B	0	0	1	∞
N	0	∞	1	> 0

1. *The GR limit: Schwarzschild solution*

To help to interpret the general case, we consider first the pure GR limit $c_1 = 0$, for which $Y_+ = -1$ and $Y_- = -\infty$. The solution is then

$$B = 1 + Y \tag{36}$$

$$N = 1/(1 + Y) \tag{37}$$

$$r_{\min}/r = Y/(1 + Y). \tag{38}$$

This is just the Schwarzschild solution, with $Y = 1/(r - r_{\min})$ and $r_{\min} = 2M$. Spatial infinity corresponds to $Y = 0$, and as $Y \to \infty$ the radius decreases to $r_{\min}$ at the bifurcation surface of the horizon. The other side of the wormhole is here labelled by the same values of Y. The range $-\infty < Y < -1$ corresponds to the future wedge of the black hole interior, where the Killing vector is spacelike. The remaining range $-1 < Y < 0$ is also significant.

It corresponds to the negative mass Schwarzschild solution.

2. Static aether solutions for generic c_1

For generic values of c_1 the limit $Y \to 0$ still corresponds to an asymptotically flat spatial infinity, where the limiting form of the solution is

$$B = 1 + Y + \cdots \tag{39}$$

$$N = 1 - Y + \cdots \tag{40}$$

$$Y = 2M/r + \cdots , \tag{41}$$

and the mass M is related to the minimum radius by

$$r_{\rm min}/2M = (-Y_+)^{-1}(-1 - Y_+)^{(1+Y_+)/(2+Y_+)}. \tag{42}$$

This ratio grows smoothly from 1 for $c_1 = 0$, to about 1.23 for $c_1 = 1$, and reaches $e/2 \approx 1.4$ for $c_1 = 2$.

Series solution in powers of $x = 2M/r$ yields

$$B = 1 + x + (1 + \nu)x^2 + \cdots \tag{43}$$

$$N = 1 - x - (\nu/6)x^3 + \cdots \tag{44}$$

$$Y = x + x^2 + (1 + \nu/2)x^3 + \cdots . \tag{45}$$

To leading order in $1/r$ this agrees with the Schwarzschild solution, as already seen previously in [13], where the two Eddington-Robertson-Schiff PPN parameters γ and β were found to agree with the GR value of 1. This completes our characterization of the asymptotically flat region. What happens when we follow the solution to smaller values of r?

The answer depends on the range of Y considered. For $Y \in (Y_+, 0)$, equation (35) or its graph in Fig. 1 indicate that $r_{\rm min}$ must be *negative*, which according to (42) implies a negative total mass $M < 0$. In this case there is a naked singularity at $Y = Y_+$ ($r = 0$) connected to an asymptotically flat region at $Y = 0$ ($r = \infty$), like in the negative mass Schwarzschild solution.

For positive Y the solution is different. It is seen again from (29) or its graph that Y grows monotonically as r decreases. As $Y \to \infty$ the r.h.s. of (35) goes to 1, so this limit for Y corresponds to a minimum radius $r_{\rm min}$, just as in the case of the Schwarzschild

solution. However, the solution behaves quite differently from Schwarzschild. First, instead of $N(r_{\min}) = 0$ we have

$$N(r_{\min}) = \left(\frac{Y_+}{Y_-}\right)^{-Y_+/(2+Y_+)}, \tag{46}$$

so the minimal 2-sphere does not sit at a Killing horizon. The value of $N(r_{\min})$ grows smoothly from 0 for $c_1 = 0$ to about 0.083 for $c_1 = 1$ and reaches $e^{-2} \approx 0.135$ for $c_1 = 2$. (Recall that in the GR limit we have $Y_- \to -\infty$.) Another difference due to the finiteness of Y_- is that the solution continues with negative Y values, with the two values $Y = \pm\infty$ identified. According to (29), as Y grows from $-\infty$ up to Y_-, r increases from $r_{\min}$ to ∞. Therefore the "interior" of the minimal 2-sphere flares out to infinite radius as in the Schwarzschild solution. But unlike the Schwarzschild case, now two values of Y correspond to each r, and the "interior" geometry is not equivalent to the exterior. In fact the difference is quite dramatic: at the internal infinity both N and B go to zero, whereas they both approach one in the asymptotically flat region.

The Carter-Penrose diagram for this solution is the square diamond in Fig. 2 with asymptotically flat past and future null infinity on the lower and upper edges bounding the right hand side and a singularity on both edges $S^{\pm}$ and at S^0 bounding the left hand side. $S^{\pm}$ are singular Killing horizons with vanishing surface gravity. The proper distance to S^0 along a constant t surface is finite if $0 < c_1 < 3/2$ and infinite if $3/2 < c_1 < 2$, while for any c_1 the affine parameter to $S^{\pm}$ along a radial light ray is finite, as we now demonstrate.

As the internal infinity at Y_- is approached, the proper radial distance on a constant t surface behaves as

$$\frac{dl}{dr} = B^{1/2} \sim (Y_- - Y)^{1/2}. \tag{47}$$

On the other hand (35) shows that in this limit the relation between Y and r is

$$(Y_- - Y) \sim r^{\frac{2+Y_+}{1+Y_+}}, \tag{48}$$

so we have

$$\frac{dl}{dr} = r^{\frac{2+Y_+}{2+2Y_+}}. \tag{49}$$

The exponent of r in (49) is always negative, and it is equal to -1 when $Y_+ = -4/3$, which corresponds to $\nu = 3/16$, i.e. $c_1 = 3/2$. For $c_1 < 3/2$ the radial distance to S^0 is finite and there is a curvature singularity at $r = \infty$ that shows up, for example, in the square of the Riemann tensor.



FIG. 2: Carter-Penrose diagram of the static aether solution. The left hand edge corresponds to spheres of infinite radius and is singular.

Along a radial null geodesic the quantity $N\dot{t} = Ndt/d\lambda$ is conserved, where λ is an affine parameter. Together with the lightlike condition $N\dot{t}^2 - B\dot{r}^2 = 0$ this implies that as the internal infinity is approached the affine parameter behaves as

$$d\lambda/dr = (NB)^{1/2} \sim (Y_- - Y)^{1/(2+Y_+)} \sim r^{1/(1+Y_+)}. \tag{50}$$

The affine parameter distance to $S^\pm$ is therefore finite for all $Y_+ \in (-2, -1)$, corresponding to all $c_1 \in (0, 2)$.

Note that since the minimal 2-sphere is not hidden by a horizon, a spherical congruence of null rays will converge towards the minimal sphere and exit the other side with a positive expansion. The Raychaudhuri equation shows that this can happen only if $R_{ab}k^a k^b < 0$ somewhere along the congruence, so we infer that the aether stress tensor must violate the null energy condition in this solution. We computed the curvature for this solution and found that $G_{tt} = -\nu Y^2/Br^2$, so the energy density of the aether ($\propto G_{tt}$) is negative *everywhere* . The solution nevertheless has positive total mass, which may at first seem to be inconsistent but it is not. The total mass of an asymptotically flat spacetime is given by a surface integral at spatial infinity, which for stationary spacetimes is proportional to

the volume integral $\int_\Sigma R_{ab} n^a \xi^b dV$, plus a surface term if there is an inner boundary [24]. Since $R_{ab} \propto T_{ab} - (1/2) T g_{ab}$, it is not the energy density that figures in the total energy but rather R_{tt}. Quite surprisingly, it turns out that R_{tt} vanishes everywhere in the static aether solution. (The only nonzero component of the Ricci tensor is R_{rr}.) Hence the energy integrand vanishes identically, as in Schwarzschild spacetime. This does not mean the total energy vanishes however, since there is a contribution from the inner boundary. In Schwarzschild that inner boundary may be pushed off to the asymptotic region on the other side of the Einstein-Rosen bridge, but the static aether solution is singular on the other side of the throat. One can think of the mass as determined by a boundary condition at this singularity.

Let us briefly consider solutions for c_1 in the range $3/2 < c_1 < 2$, corresponding to $3/16 < \nu < 1/4$. In this case the distance to the internal infinite radius sphere is infinite, and all the algebraic and differential invariants that we checked (including R, $R_{ab}R^{ab}$, $R_{abcd}R^{abcd}$, $R_{ab}u^a u^b$, and $(\nabla_a u^b)(\nabla_b u^a)$) are asymptotically zero. However, the curvature component $R_{ab}k^a k^b$ blows up asymptotically, when k^a is the tangent to an affinely parameterized radial null geodesic approaching the internal infinity. The invariant $u^a k_a$ blows up as $N^{-1/2}$, since $N^{1/2}u^a$ is the Killing vector ξ^a and $k^a \xi_a$ is conserved along the geodesic. This suggests that the above-mentioned invariants vanish because the tensor structure of the curvature, the aether, and all derivatives is determined by a single null vector pointing in the future radial null direction opposite to k^a, i.e. pointing away from the internal area-infinity.

Returning now to the generic solution for $0 < c_1 < 1$, we examine more closely the behavior at the throat and at the internal infinity. Since the spherical radius r is not a good coordinate at the minimal area sphere, nor at the internal area-infinity, we adopt instead the proper length coordinate l, in terms of which the line element takes the form

$$ds^2 = N(l)dt^2 - dl^2 - r(l)^2 d\Omega^2 \tag{51}$$

To get an idea of how the throat geometry depends on c_1, we fix the mass M of the solution and plot in Fig. 3 the numerically computed function $r(l)$ for several different values of c_1. There is a discontinuity at $c_1 = 0$ where the solution abruptly changes from a singular flare-out in finite proper distance to a perfectly regular Einstein-Rosen bridge. The singularity approaches the throat as $c_1 \to 0$, but in the same limit the curvature becomes finite and the other half of the bridge suddenly appears.



FIG. 3: Plot of area radius r vs. proper length l for fixed mass M, in units with $2M = 1$, for $c_1 = 0$, 0.1, 0.7, and 1.9. In the GR case $c_1 = 0$ this is the Einstein-Rosen bridge. For $c_1 = 0.1$ the radius flares out to infinity so quickly that the code used to make the plot halted at small radius. With increasing c_1 the throat widens, the flare-out inside is slower, and the proper length to the curvature singularity increases, becoming infinite for $c_1 \geq 3/2$.

To more fully compare the Schwarzschild and aether solutions we plot together in Fig. 4 the radius $r(l)$ and the norm of the Killing vector $\sqrt{N(l)}$ for the two solutions with the same value of the total mass M. At the internal singularity the norm of the Killing vector goes to zero, and the Killing vector is tangent to the constant r surfaces, so the singularity is a "would-be" Killing horizon. The surface gravity of the horizon is given by $d\sqrt{N}/dl$ at the horizon. The behavior is easily found using the previous formulae:

$$d\sqrt{N}/dl = (1/2)(N/B)^{1/2}(Y/r) \propto (Y - Y_-)^{2(1+Y_+)/(2+Y_+)}. \tag{52}$$

The exponent is positive, so the derivative of the norm vanishes as $Y \to Y_-$. The surface gravity is therefore zero, so the would-be Killing horizon is extremal.



FIG. 4: Plot of $r(l)$ and the norm of the Killing vector $\sqrt{N(l)}$ for GR and for $c_1 = 0.5$, for the solution with the same value of the total mass M, in units with $2M = 1$. In GR N vanishes at the bifurcation sphere at the center of the Einstein-Rosen bridge. In the ae-theory solution the Killing vector remains timelike at the throat, but at the internal $r = \infty$ curvature singularity both the norm and its slope vanish, indicating the presence of a singular extremal Killing horizon.

3. Charged dust interpretation

We argued at the beginning of this section that the c_2 and c_3 terms in the action do not contribute to the field equations in the case of static, spherical symmetry and static aether, and the c_4 term can be absorbed into a simultaneous shift of c_1 and c_3. This enabled us to reduce the general case to the one with only c_1 non-zero. We can nevertheless choose to include a c_3 term in the action, and by so doing re-express the content of the field equations in an interesting way. In particular, if we choose $c_3 = -c_1$, then the c_1 and c_3 terms combine to make $(c_1/2)F_{ab}F^{ab}$, where $F_{ab} = \nabla_a u_b - \nabla_b u_a$. This is just the Maxwell Lagrangian for a vector potential u_a, up to a constant factor. We have been treating the contravariant vector

u^a as the independent field variable, but in this Maxwell-like case it is natural to adopt instead the covariant vector u_a as independent. This change just amounts to an invertible field redefinition, hence yields the same equations of motion when the metric and aether field equations are taken together. With this choice of field variable the theory with $c_3 = -c_1$ and $c_2 = c_4 = 0$ looks quite similar to Maxwell theory, the only difference being the constraint term $\lambda(g^{ab}u_a u_b - 1)$ in the Lagrangian. It was shown in [6] that this is equivalent to the Einstein-Maxwell-charged dust system, with a constant charge to mass ratio fixed by c_1, and restricted to the sector in which there exists a gauge choice for which the vector potential is parallel to the dust 4-velocity. (This is a real restriction.)

It must be possible to interpret our strange static aether wormhole solution as a charged dust solution, but it is at first hard to see what could stabilize the dust unless it is extremally charged, which corresponds to the case $c_1 = 2$. In fact it was argued in Ref. [6], invoking prior results [25] for the charged dust problem, that there is no static solution with static aether and general c_1. However, our result here shows that argument cannot be correct. The contradiction is resolved by the observation that the prior results invoked in Ref. [6] apply only if the dust mass density is positive, whereas in the solution at hand this mass density turns out to be negative. (In Ref. [26] the charged dust system with negative mass density and constant charge to mass ratio was studied (among other cases) in static axisymmetry, and it was shown that every harmonic function determines a solution. Presumably among these solutions is the static aether solution found here.) This is related to the negative energy density that we already inferred above must be present. With a negative mass density, the dust is gravitationally repulsive, and though it has the same sign charge it is electrically attractive. (In Newtonian terms, a force $\mathbf{F}$ produces an acceleration $\mathbf{F}/m$ which is opposite to $\mathbf{F}$ when m is negative.) Thus the gravitational and electric forces exchange their usual roles. The fact that the dust does not then just collapse on itself is perhaps due to the associated cost in (positive) electric field energy when the dust is squeezed together. Although static, the solution is not regular, since there is an internal singularity.

4. Solutions with $Y \in (Y_-, Y_+)$

So far we have discussed the solution for Y in all ranges except for $Y \in (Y_-, Y_+)$. In this range the metric function B is negative, so ∂_r is timelike. Therefore, in order for the metric

signature to be Lorentzian, ∂_t must be spacelike. But in this static solution the aether is parallel to ∂_t (cf. (16)), so cannot be timelike. Hence there is no Lorentzian solution with timelike aether corresponding to this range of Y.

As a mathematical curiosity, if we allow $N > 0$, so there are two timelike dimensions, there would still be a further restriction for a real solution, since the ratio in (34) is negative and raised to the power $-Y_+/(2+Y_+)$. In order for N to be real and positive this power must be an even integer m, which implies $\nu = (1/4)(1-1/m^2)$. The ratio in (35) is also negative, and is raised to the different power $1/(2+Y_+)$, which can not also be an integer since Y_+ is not an integer. However, the integration constant $r_{\rm min}$ can be complex, thus balancing the phase of the right hand side of (35), and yielding a real solution with signature $(++--)$.

V. STARS

In this section we investigate the static solutions to the theory that, unlike the singular wormhole have a regular origin. We first show that there are no such regular pure aether solutions, and then turn to the case of fluid stars.

A. Nonexistence of pure aether stars

Spherically symmetric self-gravitating "solitons" appear in a number of field systems coupled to gravity. For example there are boson star solutions in the Einstein-Klein-Gordon system [27] and Einstein-Yang-Mills theory possesses the Bartnik-McKinnon solutions [28, 29]. It is therefore natural to ask whether such"aether star" solutions might exist in the vacuum Einstein-Aether theory. The static aether solution studied in the previous section is the unique solution with static aether (remember, this means aether aligned with the timelike Killing vector), and does not have a regular origin. Thus the only way a regular aether star might exist is if the aether has a radial component. We now examine this possibility and show that it cannot occur.

The analysis of Section III showed that local solutions around a general $r = r_0$ are characterized by three free parameters which may be taken to be $A'(r_0)$, $a(r_0)$ and $a'(r_0)$. If we apply this result at the origin $r_0 = 0$ the parameter freedom is restricted. Spherical symmetry implies that at the origin the radial component of the aether vanishes, $b(0) = 0$.

The normalization constraint (12) therefore fixes $a(0) = e^{-A_0/2}$, and the r-derivative of this constraint implies that $A'(0)$ and $a'(0)$ are not independent, but rather are related by $A'(0) = -2a'(0)/a(0)$. Thus there is a one parameter family of solutions regular at the origin.

These solutions cannot be asymptotically flat, for the following reason. The asymptotically flat boundary condition discussed in Section III would require fixing the one free parameter, leaving a unique solution. However, pure ae-theory is scale invariant, so there must be at least a one parameter family of solutions much like the Schwarzschild solutions of different mass in GR. (By contrast, Einstein-Yang-Mills theory is *not* scale invariant, and admits a discrete family of soliton and black hole solutions.) We conclude that no regular aether stars exist. This conclusion was verified empirically by integrating out from the origin with different initial parameters, and attempting unsuccessfully to tune to an asymptotically flat solution.

B. Fluid Stars

Although there are no regular vacuum aether stars, globally regular solutions exist in the presence of a static, spherically symmetric perfect fluid with no aether couplings. The fluid stress tensor appearing in the metric field equation (5) is

$$T^M_{ab} = \big(\rho(r) + P(r)\big)v_a v_b - P(r)g_{ab} \tag{53}$$

where $v^a = e^{-A/2}(\partial_t)^a$ is the fluid 4-velocity, $\rho(r)$ it's mass density, and $P(r)$ it's pressure. The metric field equation and the Bianchi identity together imply that the sum of the aether and fluid energy-momentum tensors is divergenceless. In addition, since the aether does not couple directly to the fluid, its stress tensor is independently divergenceless when its field equation and unit constraint are satisfied. Therefore the fluid stress tensor is also independently divergenceless in any solution. Thus, an appropriate system of equations for the aether plus fluid case is the (i) metric field equation, (ii) aether field equation, (iii) radial component of $\nabla^a T^M_{ab} = 0$, which is the hydrostatic equilibrium equation for the fluid

$$P' + \tfrac{1}{2}A'(\rho + P) = 0, \tag{54}$$

and (iv) an equation of state $\rho = \rho(P)$.

The count of parameters that can be freely specified when integrating out from a regular origin is the same as for the vacuum aether solution, except for the addition of a central value for the pressure $P_0 = P(0)$. For a fixed central pressure, there is just one parameter which can be tuned to obtain an asymptotically flat solution. Now there is no contradiction with scale invariance, since the central pressure sets the scale and determines the total mass of the solution. That these asymptotically flat star solutions have a static aether can be inferred as follows. The field equations can be integrated out from the origin with the aether assumed static. The pressure drops to zero at some value $r = R$, the surface of the star, where the static interior solution can be matched to the static vacuum aether solution discussed in section IV. This solution is asymptotically flat, so it must be the unique asymptotically flat solution whose existence is indicated by the parameter count.

A possible worry about the preceding argument is whether the matching to a vacuum solution at the surface of the star could be non-unique. Provided $a(r)$, $A(r)$, $B(r)$ and their derivatives are continuous at $r = R$ one can simply continue the integration into the vacuum region. We have not studied this behavior in detail, but it seems that as long as this timelike surface is not a characteristic surface of the ODE's corresponding to a spherical wavefront of the spin-0 modes of the theory, the field equations will imply the continuity condition.

To determine a star solution one can fix a central pressure and numerically integrate the tt and rr metric field equations and the hydrostatic equilibrium equation (54), using the equation of state, from the origin to the radius R where the pressure vanishes. There $A'(r)$ is continuous so one can use it to match to the vacuum solution. The total mass M can be read off from (42) together with (35), using the definition (27), $Y(R) = RA'(R)$. The area of the 2-spheres in such a star solution is strictly increasing as r increases from zero to the surface of the star where P vanishes. At that point $P' \leq 0$, so according to (54) $A' \geq 0$ (assuming positive fluid energy density ρ). Thus (27) implies $Y \geq 0$, which means that we always match to the static aether solution *outside* of the minimal area 2-sphere. A "throat" never occurs in such a star solution.

1. Constant density stars

To get a sense of the nature of the static aether star solutions we consider here the simplest example, stars with constant energy density interior. Although this does not closely describe



FIG. 5: Total mass vs. central pressure P_0 in a constant density star, in units with $\rho = 1$ and $8\pi G = 1$, for several values of c_1. The GR curve asymptotically approaches $M = (4\pi/3)(24/9)^{3/2}$. As c_1 grows the maximum mass decreases, and the curve develops a sharp local maximum and a shallow local minimum.

realistic stars, it is adequate for indicating the behavior of maximum mass limits and the stability properties of equilibrium configurations. The discontinuity in the mass density at the surface entails via the field equations a jump in A'', but A' remains continuous so can be used to match to the vacuum solution as described above.

Graphs of M versus the central pressure P_0 for the equilibrium configurations are displayed in Figure 5. In GR the mass asymptotes to a maximum value as the central pressure goes to infinity. Physically, an infinite central pressure would be required to maintain equilibrium for a greater mass. As c_1 increases in the Einstein-Aether case the maximum mass limit decreases, and the mass curve develops a local maximum and a very shallow local minimum that is only apparent for larger values of c_1. For sufficiently large c_1 a second local maximum occurs. (We have not attempted to determine the behavior at arbitrarily high pressures and for c_1 approaching 2. Perhaps the series of maxima and minima continues.)



FIG. 6: Total mass vs. R for a constant density star, in units with $\rho = 1$ and $8\pi G = 1$, for P_0 up to 300 and several values of c_1. For small pressures all the curves increase uniformly. For $c_1 = 0.005$ the slope is nearly the same as in GR. For $c_1 = 0.05$ a maximum occurs for a large central pressure. By $c_1 = 0.6$, the first maximum occurs at much smaller pressures, and there is also a minimum. For $c_1 = 1.6$ a second maximum has appeared.

The presence of stationary points in the mass versus pressure curves is an indication that the stability character of the equilibrium may be changing [30]. The connection with linearized stability arises as follows. If the squared frequency of a mode is positive then the corresponding perturbation of the star is oscillatory, while if it is negative the perturbation grows exponentially in time. In the borderline case of zero frequency the mode has zero energy, hence corresponds to a variation between two static solutions with the same mass. Transitions between stability and instability therefore occur at extrema of the mass versus R plot shown in Figure 6, where a small displacement of R does not change the mass to first order.

In the GR limit there is no critical point. The mass increases monotonically with central pressure, as seen in Fig. 5, so there is no onset of instability. In ae-theory even flat space is

not necessarily stable. The conditions on the c_i for which all linearized plane wave modes have positive squared frequency were found in [16]. For example in the pure c_1 case they are $-2 < c_1 < 1$. If we assume the values of c_i are such that very small mass stars are stable, then instability can only set in at a critical point of the mass function. As c_1 grows larger, the curves in Fig. 6 exhibit extremal points corresponding to the local maxima and minima of Fig. 5. At the maximum mass the lowest mode becomes unstable. At the following local minimum another zero frequency mode occurs, corresponding to the next mode becoming unstable. (It cannot be the lowest mode becoming stable again, since R is increasing with increasing central pressure, implying the presence of a node in the corresponding radial mode [30].) Therefore, beyond the maxima in shown in Figure 6 constant mass density stars in the Einstein-Aether theory are unstable. For small values of c_1 the central pressure has to be very large compared to the density for the star to reach the instability, implying a violation of the dominant energy condition. For $c_1 = 1$ the pressure at the onset of instability is about 1.28 times the density.

VI. DISCUSSION

In this paper we analyzed the Einstein-Aether theory assuming stationary spherical symmetry. We determined the number of free parameters in the corresponding solutions to the field equations, and classified the asymptotically flat ones. The vacuum solution with static aether, i.e. aether aligned with the timelike Killing vector, was found analytically up to inversion of a transcendental equation. It has a wormhole-like structure, with a minimal 2-sphere and a singular internal area-infinity which lies at finite affine parameter along a radial null geodesic and finite proper distance if $c_1 < 3/2$. The static fluid star solutions were classified and studied numerically for the case of constant density, and the maximum mass and stability properties were determined.

Several directions for further work are suggested by these results. Foremost from an observational point of view might be to determine the maximum mass limits for a realistic neutron star equation of state. For astrophysical applications it would also be necessary to determine the structure of rotating solutions. Another open issue is the condition on the c_i required by stability of small mass stars. A related question is whether the pure static aether wormhole solution is stable.

The solutions studied here may be of some help on the unresolved question of energy positivity in ae-theory. It is known for what ranges of the coefficients c_i the energy of linearized solutions is positive [17, 19], and it is known that for the Maxwell-like special case ($c_3 = -c_1$, $c_2 = c_4 = 0$) nonsingular negative energy initial data exist [12, 31]. The positive mass static aether solutions on the other hand have positive energy despite having everywhere negative energy density and an interior singularity, as explained in section IV B 2. It would be interesting to see whether a positive energy result can be established for nonsingular static, spherically symmetric star solutions.

Finally, we found that there is a two parameter family of static, asymptotically flat vacuum solutions. Stellar exteriors for different masses form a one-parameter family. The other parameter pertains to the radial tipping of the aether away from the Killing vector. In a black hole solution the aether must tip, since it cannot remain timelike and be aligned with the null Killing vector on the horizon. In the companion to this paper [22] we determine the structure of black hole solutions in Einstein-Aether theory.

Acknowledgements

This research was supported in part by the NSF under grant PHY-0300710.

---

[1] D. Mattingly, "Modern tests of Lorentz invariance," Living Rev. Rel. **8**, 5 (2005) [arXiv:gr-qc/0502097].

[2] C.M. Will and K. Nordtvedt, Jr., "Conservation Laws and Preferred Frames in Relativistic Gravity. I. Preferred-Frame Theories and an Extended PPN Formalism," Astrophys. J. **177**, 757 (1972); K. Nordtvedt, Jr. and C.M. Will, "Conservation Laws and Preferred Frames in Relativistic Gravity. II. Experimental Evidence to Rule Out Preferred-Frame Theories of Gravity," Astrophys. J. **177**, 775 (1972); R.W. Hellings and K. Nordtvedt, Jr., "Vector-metric theory of gravity," Phys. Rev. **D7**, 3593 (1973).

[3] See, for example, M. Gasperini, "Classical Repulsive Gravity And Broken Lorentz Symmetry," Phys. Rev. D **34**, 2260 (1986); "Singularity prevention and broken Lorentz symmetry", Class. Quantum Grav. **4**, 485 (1987); "Repulsive gravity in the very early Universe", Gen. Rel.

Grav. **30**, 1703 (1998); and references therein.

[4] V. A. Kostelecky and S. Samuel, "Gravitational Phenomenology In Higher Dimensional Theories And Strings," Phys. Rev. D **40**, 1886 (1989).

[5] M. A. Clayton and J. W. Moffat, "Scalar-Tensor Gravity Theory For Dynamical Light Velocity," Phys. Lett. B **477**, 269 (2000) [arXiv:gr-qc/9910112].

[6] T. Jacobson and D. Mattingly, "Gravity with a dynamical preferred frame," Phys. Rev. D **64**, 024028 (2001) [arXiv:gr-qc/0007031].

[7] N. Arkani-Hamed, H. C. Cheng, M. A. Luty and S. Mukohyama, "Ghost condensation and a consistent infrared modification of gravity," JHEP **0405**, 074 (2004) [arXiv:hep-th/0312099].

[8] B. M. Gripaios, "Modified gravity via spontaneous symmetry breaking," JHEP **0410**, 069 (2004) [arXiv:hep-th/0408127].

[9] R. Bluhm and V. A. Kostelecky, "Spontaneous Lorentz violation, Nambu-Goldstone modes, and gravity," Phys. Rev. D **71**, 065008 (2005) [arXiv:hep-th/0412320].

[10] C. Heinicke, P. Baekler and F. W. Hehl, "Einstein-aether theory, violation of Lorentz invariance, and metric-affine gravity," Phys. Rev. D **72**, 025012 (2005) [arXiv:gr-qc/0504005].

[11] V. A. Rubakov, "Phantom without UV pathology," arXiv:hep-th/0604153.

[12] C. Eling, T. Jacobson and D. Mattingly, "Einstein-aether theory," in *Deserfest*, eds. J. Liu, M. J. Duff, K. Stelle, and R. P. Woodard (World Scientific, 2006) arXiv:gr-qc/0410001.

[13] C. Eling and T. Jacobson, "Static post-Newtonian equivalence of GR and gravity with a dynamical preferred frame," Phys. Rev. D **69**, 064005 (2004) [arXiv:gr-qc/0310044].

[14] M. L. Graesser, A. Jenkins and M. B. Wise, "Spontaneous Lorentz violation and the long-range gravitational preferred-frame effect," Phys. Lett. B **613**, 5 (2005) [arXiv:hep-th/0501223].

[15] B. Z. Foster and T. Jacobson, "Post-Newtonian parameters and constraints on Einstein-aether theory," Phys. Rev. D **73**, 064015 (2006) [arXiv:gr-qc/0509083].

[16] T. Jacobson and D. Mattingly, "Einstein–Aether waves," Phys. Rev. D **70**, 024003 (2004) [arXiv:gr-qc/0402005].

[17] E. A. Lim, "Can We See Lorentz-Violating Vector Fields in the CMB?," Phys. Rev. D **71**, 063504 (2005) [arXiv:astro-ph/0407437].

[18] J. W. Elliott, G. D. Moore and H. Stoica, "Constraining the new aether: Gravitational Cherenkov radiation," JHEP **0508**, 066 (2005) [arXiv:hep-ph/0505211].

[19] C. Eling, "Energy in the Einstein-aether theory," Phys. Rev. D **73**, 084026 (2006)

[arXiv:gr-qc/0507059].

[20] S. M. Carroll and E. A. Lim, "Lorentz-violating vector fields slow the universe down," Phys. Rev. D **70**, 123525 (2004) [arXiv:hep-th/0407149].

[21] B. Z. Foster, "Radiation damping in Einstein-aether theory," Phys. Rev. D **73**, 104012 (2006) [arXiv:gr-qc/0602004].

[22] C. Eling and T. Jacobson, "Black holes in Einstein-aether theory," Class. Quantum Grav., to appear, [arXiv:gr-qc/0604088].

[23] B. Z. Foster, "Metric redefinitions in Einstein-aether theory," Phys. Rev. D **72**, 044017 (2005) [arXiv:gr-qc/0502066].

[24] R. M. Wald, *General Relativity*, (University of Chicago Press, 1984).

[25] U. K. De and A. K. Raychaudhuri, "Static Distribution of Charged Dust in General Relativity," Proc. Royal Soc. A **303** 47 (1968).

[26] W. B. Bonnor, "Equilibrium of charged dust in general relativity," Gen. Rel. Grav. **12** 453 (1980).

[27] P. Jetzer, "Boson stars," Phys. Rept. **220**, 163 (1992).

[28] R. Bartnik and J. Mckinnon, "Particle - Like Solutions Of The Einstein Yang-Mills Equations," Phys. Rev. Lett. **61**, 141 (1988).

[29] M. S. Volkov and D. V. Gal'tsov, "Gravitating non-Abelian solitons and black holes with Yang-Mills fields," Phys. Rept. **319**, 1 (1999) [arXiv:hep-th/9810070].

[30] J. B. Hartle, *Gravity: An Introduction to Einstein's General Relativity*, (Addison Wesley, 2002), Chapter 22.

[31] M. A. Clayton, "Causality, shocks and instabilities in vector field models of Lorentz symmetry breaking," arXiv:gr-qc/0104103.